MATTHEW GIBNEY CATHOLIC PRIMARY SCHOOL STUDENT ASTHMA RECORD

Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.		sthma Manag			
Medication requirements: Name of Medication Method (Puffer or Spacer) When and how much? In an Emergency follow the Plan below that has been ticked (✓) Standard Asthma First Aid Plan Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is still little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	_				
Name of Medication Method (Puffer or Spacer) When and how much? In an Emergency follow the Plan below that has been ticked (✓) Standard Asthma First Aid Plan Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the student take 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.			·		
In an Emergency follow the Plan below that has been ticked (✓) Standard Asthma First Aid Plan Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.		<u>-</u>	_	When and how much?	
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Name of	Wedication	Method (Fuller of Spacer)	When and now much?	
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.					
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.					
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.					
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	In an Fm e	eraency follow th	e Plan below that has been ti	cked (✓)	
Step 1 Sit the student upright, remain calm and provide reassurance. Do n leave student alone. Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.				ched (*)	
Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Standar	d Asthma Firs	st Aid Plan		
Step 2 Give 4 puffs of a blue reliever puffer (Airomir, Asmol, Bricanyl or Ventolin) puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Step 1	Sit the student upright, remain calm and provide reassurance. Do not			
puff at a time, preferably through a spacer device *. Ask the studentake 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	-	leave student	alone.		
take 4 breaths from the spacer after each puff. Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Step 2	Give 4 puffs of	f a blue reliever puffer (Airomin	r, Asmol, Bricanyl or Ventolin), Or	
Step 3 Wait 4 minutes Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.					
Step 4 If there is little or no improvement, repeat steps 2 and 3. If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.		take 4 breatils	s nom the spacer after each p		
If there is still little or no improvement, call an ambulance immediate (Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Step 3	Wait 4 minutes			
(Dial 000) Continue to repeat steps 2 and 3 while waiting for the ambulance.	Step 4	If there is little	or no improvement, repeat s	teps 2 and 3.	
ambulance.				•	
		` '	intinue to repeat steps 2 and	3 while waiting for the	
* Note: Like a blue reliever puffer on ite own if no epacer available	* Note:	Lico a blue reliev	Use a blue reliever puffer on its own if no spacer available.		
Ose a blue reliever puller off its own if no spacer available.	Note.	oce a blac relievel parter of its own if its spacer available.			
	My Chile	d's Asthma Fi	rst Aid Plan (attached)		
My Child's Asthma First Aid Plan (attached)	Additions	al Commente:			
	Additions	ai Comments.			
My Child's Asthma First Aid Plan (attached) Additional Comments:					
Additional Comments: I authorise school staff to follow the preferred Asthma First Aid Plan and assist my child with	these instru	ctions. Please conta	act me if my child requires emerger		
Additional Comments:	Cianat	of Doront/Comm		Data	
Additional Comments: I authorise school staff to follow the preferred Asthma First Aid Plan and assist my child with asthma medication should they require help. I will notify you in writing if there are any chan these instructions. Please contact me if my child requires emergency treatment or if my child requires.	oignature	or Parent/Carer:		Date	